

B

(B) 4 terms

2 A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

CreatorOrder ByDate

Owned All Term (Ascending) Term (Descending) Creation date (Ascending) Creation date (Descending) Creator (Ascending)

Search

Creator (Descending) Term Definition Abbreviations Synonyms Labels

Collapse all

Creation period

Today Last week Last month Reset

From

To

Filter Cancel

Delete term

You are about to delete the term :

Confirm Cancel

Bartlett's Test Created Jun 12, 2019 (11:32) by Anurekha VIJAYARAGHAVAN

Definition

Link Link

6 sigma: Type of hypothesis test that compares variances of two or more normally distributed populations.

Label(s)

- English

Beta Risk Created Jun 12, 2019 (11:33) by Anurekha VIJAYARAGHAVAN

Definition

Link Link

6 sigma: Risk of concluding that two characteristics are the same when they are actually different.

Label(s)

- English

Bias Created Jun 12, 2019 (11:33) by Anurekha VIJAYARAGHAVAN , last modified Aug 18, 2023 (06:59) by GUENNOUNI, Nathalie

Definition

Link Link

ISO 3534-2:200: difference between the expectation of the test results and a true value

Note 1 to entry: Bias is the total systematic error as contrasted to random error. There can be one or more systematic error components contributing to the bias. A larger systematic difference from the accepted reference value is reflected by a larger bias value.

Note 2 to entry: The bias of a measuring instrument is normally estimated by averaging the error of indication over an appropriate number of repeated measurements. The error of indication is the "indication of a measuring instrument minus a true value of corresponding input quantity".

Note 3 to entry: In practice, the accepted reference value is substituted for the true value.

Label(s)

- English

Boxplot Created Jun 12, 2019 (11:33) by Anurekha VIJAYARAGHAVAN , last modified Aug 18, 2023 (06:52) by GUENNOUNI, Nathalie

Definition

Link Link

La boîte à moustache est un diagramme résumant quelques caractéristiques de position d'un jeu de données: la médiane, les quartiles, le minimum, le maximum...). Il est utilisé principalement pour comparer un même facteur dans deux jeux de données différents

Label(s)

- French